POLLUX DATABASE – CONTENT & USER'S GUIDE

Ana Palacios – Michèle Sanguillon - Stella-Maria Renucci - Alexandre Letellier

Server Version 2.2 - DataBase Version v12 – July 2024

Contents

1	AIM	2
2	SOURCES FOR THE THEORETICAL DATA 2.1 Radiative transfer and spectral synthesis codes 2.2 Atomic and Molecular data 2.3 3–D models 2.4 PHOENIX models in memoriam Dr. France ALLARD ⁺	4 4 5 5
3	THE POLLUX DATABASE - LATEST RELEASE3.1Updates of the latest release	5 6 11 13 14 16
4	 HOW TO USE 4.1 Collection Query Interface	22 22 24 24 27 28 29 33 34
5	FUTURE DEVELOPMENTS5.1ATLAS data to fill the gap of warm stellar spectra5.2Extension of the database to IR5.3(Re-)Introduction of SEDs5.4Companion data and associated services5.5Development of an interface to query and run MARCS/Turbospectrumv2.0 from the POLLUX web interface	40 40 40 40 40 40
6	CREDITS	40
7	REFERENCES	41

1 AIM

POLLUX is a database of stellar spectra developed at the Laboratoire Univers et Particules de Montpellier (LUPM - University of Montpellier - CNRS). Its aim is to provide a comprehensive library of theoretical stellar spectra with a broad coverage of the atmospheric parameters (effective temperature $T_{\rm eff}$, gravity log g and metallicity [Fe/H]) as well as spectral types across the Hertzsprung-Russell Diagram.

The POLLUX database collects and presents essentially synthetic spectra computed at high resolution (HRSS data). Some spectral energy distributions are also available only for the early spectral type models.

The HRSS are available for spectral types from O to M and for several initial chemical compositions ([Fe/H] and $[\alpha/Fe]$).

POLLUX spectra are expected to be useful to astrophysicists for stellar or galactic applications in several respects :

- abundance determinations
- accurate determination of fundamental properties of stars
- multi-wavelength coverage
- test for the current state-of-the-art model atmospheres
- stellar populations synthesis,
- as well as for teaching purposes oriented toward spectroscopy, model atmospheres, etc...

In its 12th version available as of July 2024, POLLUX is made available on-line to the community via the web page (http://pollux.oreme.org) regrouping a documentation, a retrieval interface for the data and an on-line graphic display tool. The web interface also offers the possibility to retrieve companion data to the spectra or SEDs (such as line-lists or associated model atmosphere), to access the full provenance of the spectra in various formats according to the ProvenanceDM of the IVOA, to broadcast the spectra using the SAMP protocol, and to make the convolution of portions of the spectra via the SPECONVOL VO-service (ivo://ov-gso/ssap/speconvol), thus allowing the user to simulate an observation. The data can be retrieved in formats compliant to the Virtual Observatory standards (namely FITS and XML VOTable). The POLLUX database can also be accessed via the VizieR service at Centre de Données de Strasbourg (CDS). It is a registered service of the VO ivo://ov-gso/ssap/pollux.

Table 1: Description of the different collections available in the POLLUX DB as of July 2024. The PLATO collections marked in red have a restricted access at present (see § 3).

Collection	Radiative Transfer	Spectrum Synthesis	T_{eff} [K]	Resolution 🌲	Spectral Range	Type	NLTE
AMBRE	MARCS ¹	TURBOSPECTRUM ²	[3500 - 8000]	> 150 000	VIS [†]	1D	No
BT-Dusty	PHOENIX ³	PHOENIX ³	[2000 - 6000]	> 100 000	VIS [‡] IR⊗	1D	Yes (atoms)
CMFGEN	CMFGEN ⁴	CMF-FLUX ⁴	[12020 - 63880]	> 300 000	UV; VIS; IR*	1D	Yes
CMFGEN-WR*	CMFGEN ⁴	CMF-FLUX ⁴	[33780 - 74300]	> 150 000	VIS [†]	1D	Yes
CMFGEN-OB-LMC-24	CMFGEN ⁴	CMF-FLUX ⁴	[23000 - 55000]	> 81 000	UV; VIS; IR*	1D	Yes
CMFGEN-OB-SMC-24	CMFGEN ⁴	CMF-FLUX ⁴	[23000 - 55000]	> 81 000	UV; VIS; IR*	1D	Yes
PLATO	MARCS ¹	TURBOSPECTRUM2.0 ⁵	[4502 - 7000]	> 150 000	₽SIV	1D	Yes
PLATO-INTENSITY	MARCS ¹	TURBOSPECTRUM2.0 ⁵	[4502 - 7000]	> 500 000	VIS®	1D	Yes
POPSYCLE	ATLAS12 ⁶	SYNTHE ⁶	[3800 - 140000]	> 295 000	⊖SIA	1D	No
RSG	MARCS ¹	TURBOSPECTRUM ²	[3300 - 4500]	> 150 000	IR [¢]	1D	No
STAGGER	STAGGER ⁷	OPTIM3D ⁸	[3899 - 7000]	20 000	UV; VIS; IR [‡]	3D	No
STAGGER-INTENSITY	STAGGER ⁷	OPTIM3D ⁸	[3899 - 7000]	20 000	UV; VIS; IR [‡]	3D	No
STAGGER-RVS	STAGGER ⁷	OPTIM3D ⁸	[3899 - 7000]	300 000	Gaia RVS•	3D	No
CMFGEN-SED	CMFGEN ⁴	CMF-FLUX ⁴	[12020 - 63880]	20 000	50-249 997 Å	1D	Yes
CMFGEN-WR-SED*	CMFGEN ⁴	CMF-FLUX ⁴	[20638 - 74300]	20 000	43-844 089 Å	1D	Yes
CMFGEN-OB-LMC-24-SED	CMFGEN ⁴	CMF-FLUX ⁴	[23000 - 55000]	20 000	50-249 997 Å	1D	Yes
CMFGEN-OB-SMC-24-SED	CMFGEN ⁴	CMF-FLUX ⁴	[23000 - 55000]	20 000	50-249 997 Å	1D	Yes
PLATO-SED*	MARCS ¹	I	[2500 - 8000]	20 000	1999–200 007 Å	1D	No
¹ Gustafsson et al. (1975, 2008) and Ple	z et al. (1992); see https	:://marcs.oreme.org ² Alvarez	: & Plez (1998) ³ Hau	ıschildt, Baron & A	.llard (1997)		

⁵ Gerber et al. (2023) ⁷ Magic et al. (2013); see also https://staggergrid.wordpress.com/ ⁴ Hillier & Miller (1998); see also http://kookaburra.phyast.pitt.edu/hillier/web/CMFGEN.htm ⁶ Sbordone et al. (2004); see also http://wwwuser.oats.inaf.it/castelli/sources/atlas12.html

 ‡ UV : 2000 to 3000 Å; VIS : 3000 to 12000 Å; IR : 12000 to 200000 Å • 8395 to 8905 Å $^{\odot}$ Resolution for SED given as a velocity resolution, the guoted value tep in velocity. $^{\bullet}$ Statistical samples of the model surface fluxes derived in the MARCS model calculations. Correspond to the flx data from MARCS database/ $^{\otimes}$ VIS : 3000 to 12000 Å; IR : ⁸ Chiavassa et al. (2009, 2018) [•] If the value is indicated as a minimum, it means that the resolution varies with wavelength as Δλ is constant over the entire spectral domain. \oplus 2900 to 10100 Å $^{\oplus}$ 4000 to 9000 Å * UV : 900 to 3000 Å; VIS : 3000 to 12500 Å(3000 to 12000 Å for low metallicity spectra); IR : 12000 to 25000 Å [†] 3000 to 12000 Å * Wolf-Rayet models - The temperature of reference given in the fourth column is not T_{eff} in this case but T_{\star} for these stars corresponding to the step in velocity. $^{\diamond}$ 10000 to 50000 Å 12000 to 200000 Å

2 SOURCES FOR THE THEORETICAL DATA

2.1 Radiative transfer and spectral synthesis codes

As of July 2024, the database gathers **49700** high (and medium) resolution synthetic flux and intensity spectra associated to spectral types O to M and to Wolf-Rayet stars, 23818 of which are publicly accessible, and **833** spectral energy distributions (e.g. SED) for O-type and Wolf-Rayet stars¹.

Table 1 references the radiative transfer codes used to generate the model atmospheres and the associated spectrum synthesis programs used to compute the synthetic spectra actually available in the POLLUX database. It also provides the temperature domain covered by each collection, the spectral resolution of the spectra, the spectral range provided (UV, VIS or IR), the type of radiative transfer code used to compute the model atmosphere (1–D or 3–D) and information on the NLTE treatment of the radiative transfer (yes or no).

Geometrical effects are taken into account according to the spectral type and gravity, and the HRSS available in the database are derived from both spherical and plane-parallel models accordingly.

2.2 Atomic and Molecular data

Information on the atomic and molecular data used in the computation of the spectra are given in the header attached to each spectrum (see § 3.3). Here is a description of these sources for the publicly available collections within the latest release (V12) of the database:

- For the spectra of the RSG collection, the atomic linelists are taken from the VALD database (Kupka et al. 2000), and they are complemented by specific molecular linelists (Plez, private communication) for cool stars. A link is provided in the header with the detailed linelists (atomic and molecular) used in the computations.
- For the spectra of the AMBRE collection computed by P. de Laverny within the framework of the AMBRE project (de Laverny et al. 2012), the atomic linelists are taken from the Opacity Project database (Badnell et al. 2005, Seaton 2005 and references therein), and are complemented by specific molecular linelists.
- For the spectra and SEDs of the CMFGEN-xxx collections, the atomic linelists are mainly taken from the Opacity Project database (Badnell et al. 2005, Seaton 2005 and references therein), and are occasionally complemented by specific linelists.
- For the spectra from the POPSYCLE collection, the atomic and molecular linelists are fully described in Branco et al. 2024. The atomic linelist is directly made available by the data producer via the Pollux DB interface as a companion data to these spectra (see § 3 and § 4).
- For the spectra of the STAGGER, STAGGER-RVS and STAGGER-INTENSITY collections the molecular and atomic linelists are the same as the last version of the MARCS model atmospheres (see Gustafsson et al. 2008 and Chiavassa et al. 2018, https://marcs.oreme.org).
- For the spectra of the BT-Dusty collection, multiple molecular and atomic linelists are used that can be directly found in a companion file to each spectrum accessible via an URL directly from the header file, the web interface or through a datalink for the users of the VO service

¹The PLATO-SED private collection also provides SEDs that are in fact samples of the surface flux between 1 300 and 200 000 Åthat are produced during the model atmosphere calculation.

(see below). All of the spectra currently in this collection use the Barber & Tennyson (2006) water vapour linelist. This file is the input file to the PHOENIX code originally labelled as ".spec.5" with the PHOENIX notation. Such files include all the setup information of each computation, and in particular the atomic and molecular linelists.

2.3 3-D models

Since the 9th DB release, spectra based on 3D Radiative HydroDynamic simulations of stellar atmospheres performed with the STAGGER code (Magic et al. 2013) are distributed in POLLUX. These spectra are computed with the OPTIM3D code and result from a disk integration and a temporal average (Chiavassa et al. 2018). These spectra assume zero microturbulence, as this parameter is no longer needed in 3D RHD simulations, in which velocity fields are self-consistently accounted for. They are available through three collections (see Table 1) : medium resolution power and intensity spectra (constant resolving power of $\lambda/\Delta\lambda = 20\ 000$) over a very large spectral range, from 2000 Å to 200 000 Å , high resolution spectra (constant resolving power of $\lambda/\Delta\lambda = 300\ 000$) over the narrow spectral range of the Gaia-RVS , from 8395 to 8905 Å. In addition to the flux spectra averaged over the stellar surface that were previously available in the database, the intensity spectra at different inclination angles $\mu = \cos\theta$, where θ is the angle with respect to the line of sight, are now also available in the present release².

2.4 PHOENIX models in memoriam Dr. France ALLARD⁺

Since the 10th release of the POLLUX DB we propose PHOENIX stellar spectra specifically computed for this database by F. Allard⁺. They are gathered in a collection named "BT-Dusty" which refers to models using the Barber & Tennyson (2006) water vapor linelist and in which dust is formed is equilibrium with the gas phase (maximum dust content). Such models are "valid" for Near-IR studies with $T_{eff} > 1700$ K.

Compared to the models distributed on the personal webpage of F. Allard⁺ (https://perso.ens-lyon. fr/france.allard/indexfr.html) and at the Lyon Data center for the Sciences of the Universe LyDU, the spectra available through POLLUX include both the absolute flux and **the flux normalized to the continuum**, as any other spectra in the POLLUX DB (see below). The normalised flux requires the knowledge of the continuum of the spectra, an information not available for the previous collections of PHOENIX spectra as distributed in the LyDU database and on F. Allard webpage. The continuum of the spectra in the BT-Dusty collection of the POLLUX database were computed following the procedure described in Kulenthirarajah et al. (2019), by using PHOENIX to compute the actual continuum for each spectrum. This is done by removing the discrete opacities, making any atomic and molecular data unavailable and by keeping dust and continuous opacities.

3 THE POLLUX DATABASE - LATEST RELEASE

Here we describe the data available within the database as of July 2024 in terms of actual files and coverage of the different parameters spaces. A count of the spectra available per spectral domain and collection is given in Table 3.

²Ten different inclination angles are available in addition to the averaged spectra $\mu = cos(\theta) = [1.00, 0.90, 0.80, 0.70, 0.50, 0.30, 0.20, 0.10, 0.05, 0.01].$

3.1 Updates of the latest release

• New CMFGEN collections

Four new collections of CMFGEN spectra are included (two collections of HRSS and associated SEDs). They correspond to models of O and B-type stars at the metallicities of the LMC (CMFGEN-OB-LMC-24 and CMFGEN-OB-LMC-24-SED) and the SMC (CMFGEN-OB-SMC-24 and CMFGEN-OB-SMC-24-SED). As for the other CMFGEN spectra they are available in the UV, VIS and IR domains (see Table 1).

• New POPSYCLE collection

A new collection providing spectra computed using the ATLAS12/SYNTHE suite is made available as the public outcome of a ANR funded project POPSYCLE led by Pr. A. Lançon (Strasbourg Observatory, France). These spectra are designed to represent the two main stellar populations of four galactic globular clusters at four different metallicities (NGC 1904, NGC 5904, NGC 0104, and NGC 5927, with respective metallicities [Fe/H] = -1.58, -1.29, -0.77, and -0.47 dex). For each point in the Kiehl diagram (log g vs T_{eff}), there is a spectrum with solar scaled abundances to the metallicity of the cluster considered, and a spectrum that is enriched in helium (Y = 0.3) and bears the imprint of the hot CNO-cycle nucleosynthesis ([C/Fe] = -0.3 dex, [N/Fe] = +1.2 dex, [0/Fe] = -0.05 dex, [Na/Fe] = +0.6 dex). All spectra are α -enhanced with [α /Fe] = +0.4 dex.

New PLATO private collection

From the present version of the Pollux Database, we give the opportunity for a consortium to use the POLLUX database and infrastructure to share data through among a restricted number of people for a fixed proprietary period. After this period, the data will become publicly available through the database. Within this framework, we have integrated the PLATO, PLATO-INTENSITY and PLATO-SED collections with restricted access. They contain optical spectra computed by the PLATO Science Preparation Management team under WorkPackage WP 122 100 "1-D stellar atmospheres" led by Pr. B. Plez. A dedicated user's guide and specific description is provided upon granted access to the database. These collections will become public by the launch of the PLATO mission in December 2026. In the meantime, visitors may ask for access to the collection to the Pollux management team who may grant it after validation by the PLATO Consortium.

3.2 The data : Synthetic Spectra and Spectral Energy Distributions

The high resolution synthetic spectra (hereafter HRSS) distributed consist of three columns files giving the wavelength, the absolute flux and the normalised flux in the first, second and third columns respectively.

All the spectra³ are available in the optical range between 3000 Å and 12000 Å, and for some collections additional spectral ranges (UV or IR) are also made available with a wavelength interval that may vary from one collection to the other (see Table 1).

The number of points in each spectrum is usually the same for all spectra belonging to a specific collection except for the BT-Dusty collection where each spectrum has a different number of points.

The resolution of the different spectra in the database is indicated in Table 1. Here are some specifications for the spectra that do not have a constant resolution :

³The only exception is the RSG collection, for which optical spectra are under production.



Figure 1: Synthetic spectra of the solar simulation in the spectral range 2000 – 25000 Å and for different $\mu = cos(\theta)$ inclination angles used in the computations of the spectra from the STAGGER-INTENSITY collections. *From Chiavassa et al. 2018, A& A 611, A11.*

- For the BT-Dusty collection, the spectral resolution varies as a function of the wavelength within each spectrum, and varies from one spectrum to another depending on the temperature with $R \ge 130\ 000$.
- For the RSG and AMBRE collections, the resolution varies within the spectral domain considered, and it is characterised by a constant step in wavelength $\delta \lambda = 0.02$ Å, leading to R \geq 150 000 for these data sets.
- For the PLATO collection, the spectral resolution varies from 570000 at the blue end to 1290000 at the red end of the spectrum.

Since the 10th release of the POLLUX database, Spectral Energy Distributions for Wolf-Rayet, O and B-type stars from the CMFGEN collection are made available. The current version of the database includes new SEDs at the metallicities of the Small and Large Magellanic Clouds. The 833 SED derived from CMFGEN model atmospheres cover the following wavelength range $\lambda \in [50; 250000]$ Å with a resolution R = 10 000. The spectral energy distributions are available in the form of a two column file giving the wavelength and the absolute flux as the first and second columns respectively.

Figures 2 show the adopted nomenclature for the data in POLLUX. Each name includes information about temperature, surface gravity, metallicity, chemical composition (with different nuclides highlighted depending on the collection), the geometry of the model atmosphere and specific variable related to the model atmosphere computation (microturbulent velocity, mass loss or parameters of the wind), as well as the specification of the spectral domain it encompasses. Table 2: Collections and corresponding prefixes in the adopted nomenclature

Collection	Prefix
AMBRE	М
BT-Dusty	Р
CMFGEN, CMFGEN-OB-SMC-24, CMFGEN-OB-LMC-24	С
CMFGEN-WR	CWR
PLATO	PL
POPSYCLE	AP
RSG	М
STAGGER	0
STAGGER-RVS	G
STAGGER-INTENSITY	0



Figure 2: Name nomenclature for the CMFGEN (first upper line) AMBRE,RSG or BT-Dusty (second line), POPSYCLE (third line) and STAGGER (lower line) files. The spectra are given the extension **.spec** and the associated headers have **.spec.txt**.

Finally the size of uncompressed individual high resolution synthetic spectra in ASCII format is given in Table 3 according to the collection and the spectral domain.

Collection	Spectral Domain	Size	Number of spectra
AMBRE	VIS	15.3 MB	12927
BT-Dusty	IR	18 to 20 MB	377
BT-Dusty	VIS	14 to 22 MB	377
CMFGEN	IR	2.2 MB	309
CMFGEN	VIS	5.2 MB	309
CMFGEN	UV	1.6 MB	309
CMFGEN-OB-LMC-24	IR	1.7 MB	272
CMFGEN-OB-LMC-24	VIS	8 MB	272
CMFGEN-OB-LMC-24	UV	1.9 MB	272
CMFGEN-OB-SMC-24	IR	1.7 MB	305
CMFGEN-OB-SMC-24	VIS	8 MB	305
CMFGEN-OB-SMC-24	UV	1.9 MB	305
CMFGEN-WR	VIS	5.8 MB	11
PLATO	VIS	20 MB	1991
PLATO-INTENSITY	VIS	20 MB	23892
POPSYCLE	VIS	26 MB	1389
RSG	IR	10 MB	40
STAGGER	IR	2.1 MB	181
STAGGER	VIS	1 MB	181
STAGGER	UV	300 kB	181
STAGGER-INTENSITY	IR	2.7 MB	1791
STAGGER-INTENSITY	VIS	1.3 MB	1791
STAGGER-INTENSITY	UV	420 kB	1791
STAGGER-RVS	VIS	655 kB	122

Table 3: Typical size of uncompressed HRSS ASCII files and Number of spectra for the different collections and spectral domains.



Figure 3: Illustration of the workflow leading to the production of a synthetic spectrum (simpler case of the 1–D model atmospheres).

3.3 The Data Model : Header files

Each spectrum can be seen as the result of a workflow as shown in Fig. 3 (codes, input physics, physical parameters characterising the spectrum, ...).

A header file describing the data itself and all the relevant ingredients used within the workfow that lead to the synthetic spectrum or the SED is attached to each dataset. In addition to this information, the header file also contains a set of descriptors characterising stellar spectra which are independent of whether the data is observed or synthetic (file structure and curation⁴ information). It also provides the original ownership of the associated spectrum.

Finally, a tag describing the detailed specifications of the spectrum that are not clearly seen from the header file is introduced, and it is used to give an appreciation of the pertinence. This tag, named "specs", appears in the form of a 8 digit number to be read as described in Table 4.

⁴Curation includes all information concerning the data sets that ensures they are available for discovery and re-use in the future. Number version of the code, data producer, date of production are part of the curation information.

Digit number	Description	Possible values
1	Winds accounted for in model atmosphere	0 = No 1 = Yes 9 = Irrelevant
2	Equality of chemical compositions for model atmosphere (code1) and synthetic spectrum (code2) *	0 = No 1 = Yes 9 = Irrelevant
3	Clouds accounted for in model atmosphere•	0 = No 1 = Yes 9 = Irrelevant
4	Narrow spectral domain [†]	0 = No 1 = Yes 9 = Irrelevant
5	Dust condensation in equilibrium with gas phase ullet	0 = No 1 = Yes 9 = Irrelevant
6	Full cloud model including nucleation, growth sedimentation and mixing in addition to condensation•	0 = No 1 = Yes 9 = Irrelevant
7	Model limitation $^{\diamond}$	0 = No 1 = Yes
8	Pertinence . Indicates whether the spectrum should be used under certain restrictions or not [‡]	1 = All applications 2 = Restricted applications

Table 4: Description of the specs tag in the header file and the database

* Some MARCS HRSS from the AMBRE project have inconsistent $[\alpha/Fe]$ values between the model atmospheres and the synthetic spectra (see de Laverny et al. 2012 for details), and are flagged with pertinence = 2. They are represented as black squares in the Fig 5. • These flags concern PHOENIX models and describe the physics of the spectra in the BT-Dusty collection. [†] This presently concerns the OPTIM3D HRSS from the STAGGER-RVS collection, are available only for a narrow portion of the visible domain. These spectra were attributed pertinence = 2. ^o This relates to limitations in the model atmosphere (code1) or the synthetic spectrum (code 2). In the present version of the database, this flag is set to 1 for the coolest spectra of the AMBRE collection ($T_{eff} \in [3500 \ K; 4500 \ K]$) for which caution is caution is advised in de Laverny et al. (2012) and Gustafsson et al. (2008). [‡] All spectra that are flagged with 0 for the 2nd digit, 1 for the 4th digit or 1 for the 7th digit have been attributed pertinence = 2, meaning they are suited for restricted uses only or to be used with caution. The spectra with Y = 0.3 from the POPSYCLE collection are flag with a pertinence = 2 because they are dedicated to the specific modeling of the 2nd population stars in globular clusters.

3.4 VO compliancy

The data provided in the POLLUX database have been made compliant to the Virtual Observatory standards.

POLLUX is accessible via the Simple Spectra Access Protocol Version 1.1, the description of which can be found at this URL http://www.ivoa.net/Documents/latest/SSA.html. This means in particular that all the relevant characteristics of the data appearing in the query forms described above have an associated UCD+1 (Unified Content Descriptors, Version 1+, see http://www.ivoa.net/Documents/latest/UCD.html) that allows for interoperability within the VO.

The POLLUX database is registered in the EURO-VO registry as a service providing theoretical spectra. The query to the registry allows POLLUX data to be visible through VO tools such as CAS-SIS (http://cassis.irap.omp.eu/), VOSpec (https://www.cosmos.esa.int/web/esdc/vospec) and Aladin (http://aladin.u-strasbg.fr/).

Since the 3rd release, the HRSS files (data + header) may be retrieved in the VO compliant formats XML VOTable, XML binary VOTable and FITS (see below). These versions of the data have been generated via the TOPCAT VO tool (http://www.star.bristol.ac.uk/~mbt/topcat/).

Since the 7th release, POLLUX can be used through the VO within the framework of Science Cases. In particular, we have developed the science case SPECFLOW (http://specflow.oreme.org) that combines the query of Vizier and Simbad CDS services, the observational database Polarbase, the POLLUX database and the spectral convolution service SPECONVOL registered in the VO (ivo://ov-gso/ssap/speconvol), included in the POLLUX web interface as described above and declared in a DATALINK adhoc service.

Since this 10th release, POLLUX provides the availability of a file containing all the lines used for BT-Dusty spectra via a service declared in a DATALINK adhoc service.

POLLUX is also accessible via the Simulation Data Access Layer (SimDAL) and the Provenance Simple Access Protocol (ProvSAP). At the moment, these two protocols are not used very often because there is still no VO tool that implements them.

As of the present release (V12 of the database), the Provenance Data Model can be retrieved in various formats for each spectrum (and SED) in the database. Moreover, we offer now the possibility to directly broadcast the spectra into VO compliant tools via a SAMP protocol (see § 4).

3.5 Coverage of the parameters space

The POLLUX database includes solar metallicity data for O to M type stars. Spectra are available in the metallicity range -5.0 dex \leq [Fe/H] \leq 1.0 dex for A to K type stars (AMBRE, PLATO, POPSYCLE and STAGGER collections).

Concerning cool and hot stars, 1D HRSS with non-solar C,N,O abundances are also provided.

Table 5 summarizes the coverage of the fundamental parameters for the different collections available in POLLUX. The newly available data as of the present 12^{th} release are highlighted in red. On the other hand, Figures (4) to (7) illustrate the coverage in the Kiel diagram (T_{eff} , log g) of the spectra in different collections.

	Idule J. COVEL	авс от ше рага	THEFT SPACE OF THE LOP		1111 2020	
Collection	T_{eff} (step) [K]	logg (step)	[M/H]•(step)	[a/Fe] (step)	x	Y
AMBRE	3500 - 4900 (200) * 5000 - 8000 (250) *	0 - 5.5 (0.5)	-5.0 – 1.0 (see Fig 5)	-0.4 – 0.8 (see Fig		1
BT-Dusty	2000 – 6000 (100)	0.5 - 5.5 (0.5)	0.0	0.0	I	1
CMFGEN	12020 – 63880 (<)	2 - 4.51 (<)	-1.48 ¹ , -0.73 ¹ ,0.0 ² 0.04; 0.05	0.0	I	1
CMFGEN-OB-LMC-24	23000 - 55000 (<)	2 - 4.51 (<)	-0.40 ³	0.0	I	I
CMFGEN-OB-SMC-24	23000 - 55000 (<)	2 - 4.51 (◊)	-0.80 ³	0.0	T	I
CMFGEN-WR	33780 – 74300 (Ø)	0.0	$-1.0 - 1.0 \ (0.5)$	0.0	$0.0-0.55^\diamond$	$0.38 - 0.99^{\diamond}$
RSG	$2300 - 4500 \ (100)$	$-1.0 - 1.0 \; (0.5)$	0.0	0.0	I	
PLATO	4500 - 7000 (<)	3 – 5 (◊)	0.0	0.0	T	I
POPSYCLE	$3800 - 14000 (\diamond)$	$1 - 4.5 (\diamond)$	-1.58;-1.29;-0.77;-0.49 ⁴	+0.4	I	0.256*, 0.300*
STAGGER	3899 – 7000 (≈ 500)	1.5 - 5 (0.5)	-4.0 – 0.5 (voir Fig. 7)	0.4; 0.2; 0.0	۱ *	I
STAGGER-RVS	$3899 - 7000 (\approx 500)$	1.5 - 5 (0.5)	-2.0 – 0.5 (voir Fig. 7)	0.4; 0.2; 0.0	۱ *	1
¹ Data at $Z = 1/5 Z_{\odot}$ and in Martins & Palacios (2(Hillier (2024) ⁴ [Fe/F NGC 0104 and NGC 593 * The effective temperatu ° Irregular stepping (see $\tau = 20$, which is the relev as can be seen from Fig. (dex for lower values of [F enhanced. [*] Correspo	I Z = $1/30 Z_{\odot}$ publishe $17) ^{3} Data at Z = 0.$ I] = $-1.48, -1.29, -0.77 a$ 27 respectively. These of the step is of 200 K for T for instance Fig. (4 for C ant scale for Wolf-Raye ant scale for Wolf-Raye 6). * α -elements are e 2e(H]. • Correspondent ands to the second populance enhanced.	d in Martins & Partino d in Martins & Partino d 10,022 and Z = 0.0 and -0.48 correspond at a republisher $e_{ff} \leq 3900$ K and CMFGEN collection t stars. Irregular inhanced for substance of the first population of the clust lation latin lation lation lation lation lation lation latin latin	alacios (2020) and in Martin: 055 for SMC and LMC respe and to the metallicities of the d in Branco et al. (2024). I of 250 K for hotter models. an). $^{\circ}$ The temperatures g stepping. The spectra are co olar metallicities with [α/Fe] dlation of the clusters, with so ters, with enriched in He and	 & Palacios (2022) ctively published in e galactic globular cl Also referred a r, i.e. iven here are T_*, i.e. mputed along evolu = +0.2 dex for [Fe/F] blar scaled abundanc I with C, N, O and N 	² Data at Z = Marcolino, Bouusters NGC 190 usters NGC 190 s [Fe/H] in most temperatures at tionary tracks of f] = -0.5 dex and es and α -elemer a initial abunda	Z _{\odot} published ret, Martins & 4, NGC 5904, cases. optical depth massive stars [α/Fe] = +0.4 ths abundance nces modified

Table 6: Parameters of the solar-type spectra

Model	T _{eff}	log g	ξ	geom.	Abundances
AMBRE	5500 K	4.5	1 km.s ⁻¹	р	GAS2007 [†]
BT Dusty	5500 K	4.5	0.86 km.s ⁻¹	S	CIFIST11 [‡]
STAGGER-RVS	5530 K	4.5	irr.	3D	GAS2007

⁺Grevesse et al.(2007) [‡]Caffau et al. (2011)

3.6 Overlapping regions

PHOENIX-based collections (BT-Dusty) and MARCS-based collections (AMBRE and RSG) overlap at solar metallicity in different regions of the Kiehl diagram. It is the case of spectra for G and K spectral type dwarf stars (see Fig. 8a, and for the spectra of K and M supergiants.

The spectra may differ in these regions due to the use of different ingredients in the computation of the model atmosphere, the synthetic spectrum or both (different input parameters, different linelists, ...).

We illustrate the overlap of the BT-Dusty, AMBRE and STAGGER collections for a solar-like star at solar metallicity in Fig. 9. The parameters used in each case are recalled in Table. 6.

Figure 9 shows the comparison between BT-Dusty, AMBRE and STAGGER-RVS (for the RVS domain only) high resolution spectra normalized to the continuum in the full optical domain, in the Gaia-RVS domain as well as in the region of the H β atomic line @ 4861 Å and of the CH line @ 4300 Å.



Figure 4: Coverage of the Kiel diagram (T_{eff} - log g) plane by the HRSS associated with (*left*) CM-FGEN (blue and cyan triangles), PHOENIX (orange circles dots) and MARCS (red and magenta squares) model atmospheres, and (*right*) PHOENIX BT-Dusty model atmospheres as of March 2023 for solar metallicity. Superimposed are standard stellar evolution tracks from the Geneva code (Schaller et al. 1992) on the left, and from Baraffe et al. (2015) on the right.



Figure 5: Distribution of the AMBRE synthetic spectra grid in the atmospheric parameters and $[\alpha/Fe]$ space. Only one value of $[\alpha/Fe]$ for every [M/H] was adopted during the model atmosphere selection process. This is illustrated in the bottom right panel where the AMBRE spectra computed from MARCS models with consistent $[\alpha/Fe]$ ratios are plotted with filled circles while open circles refer to all the other AMBRE spectra computed with atmosphere models that have inconsistent $[\alpha/Fe]$ ratios. These last ones are flagged with a pertinence = 2 in the POLLUX Database *.From de Laverny et al. (2012), A& A 544, A126.*



Figure 6: Coverage of the $(T_{eff} - L)$ plane by the HRSS associated with CMFGEN model atmospheres Wolf-Rayet stars. The overplotted tracks are stellar evolution models including rotation from Meynet & Maeder (2003) for different initial masses as labelled (dotted lines for $M_{ini} = 85 M_{\odot}$, solid lines for $M_{ini} = 60 M_{\odot}$, dot-dashed lines for $M_{ini} = 40 M_{\odot}$ and dashed lines for $M_{ini} = 25 M_{\odot}$).



Figure 7: Kiel diagram showing the coverage of the STAGGER grid on which STAGGER and STAGGER-RVS collections are based. *From Magic et al. 2013, A& A 557, A26.*



Figure 8: Illustration of the overlapping region of BT-Dusty and AMBRE collections at solar metallicity.



Figure 9: Overplot of spectra from the BT-Dusty (green), AMBRE (black) and STAGGER-RVS (blue) collections at $T_{eff} = 5500$ K, log g = 4.5 and [Fe/H] = 0. The wavelength are given in the air and the ordinate presents the logarithm of the absolute fluxes.



Figure 10: Pollux homepage.

4 HOW TO USE

The POLLUX database is accessible via the URL: http://pollux.oreme.org.

The web page is designed to be user friendly, and to allow inexperienced users to easily access, visualise and retrieve the data.

The homepage allows to access the database via two methods : 1) explore the different collections which correspond to ensembles of spectra generated using the same radiative transfer and spectral synthesis codes; or 2) query the stellar parameters. This organisation is meant to facilitate the exploration and query.

From the homepage, the user can also access the news, this user's guide, read more about the people involved in the project and contact the managers via a contact form. There is a new **RESTRICTED** ACCESS entry in the upper right menu that allows the users (and managers) to connect to access private collections (the PLATO consortium collections as of July 2024). Logged users access the private collection requested in addition to the public database. Finally, direct access to the companion tools and databases SPECFLOW and CASSIS are also provided.

4.1 Collection Query Interface

Since the 11th release of the database (March 2023), the spectra and SEDs have been separated into collections associated with either a project or a publication (STAGGER-RVS, AMBRE, POPSYCLE), a type of objects (RSG, CMFGEN-WR) or a suite of radiative transfer and spectral synthesis codes (CMFGEN, BT-Dusty, STAGGER). Each collection is presented in this page (see Fig. 12) in terms of codes used, stellar parameters covered, number of spectra/SEDs available. The data producer is also emphasized and in some cases, reference papers are also given.

Login	
Email	
Password	\$
Forgot password?	

Figure 11: Connection interface opened when clicking on the RESTRICTED ACCESS text in the upper menu.

THE SYNTHETIC STELLAR SPECTRA DATABASE POLLUX		SPECTI	RA COLLECTIONS STELLAR PARAMETERS USER'S GUIDE MY SPECTRA
Sort in alphabetical order v Some collections are o	STELLAR SPECT nly accessible upon request as members of designated consortia. You may ask for access by	RA COLLECTIONS y sending a message to the e-mail address provided in the CONTACT section of the welcom	Search
AMBRE Based on MARCs model atmospheres and TURODSPECTRUM spectral symbols Vol range: Proff (5 26): 2011 (Jeans Mrd (5 26 2.8.1)) Proff (5 26): 2011 (Jeans Mrd (5 26 2.8.1)) Resultion : 1300 Proff (5 2.6.2.1) Proff (5 12.6.2.1) Proff (5 2.6.2.1) Proff (5 2.6.2.1) Proff (5 2.6.2.1) Proff (5 2.6.2.1) Description (1 2.6.2.1)	BT-Dusty Stacd on PHOCHXII: model attrinospheres VS and dis runges Myder [E] (b a. 2.8) /Pyder [C] (b a. 2.8) Resultation: 10.0000 Produced by F. Allied (CRAL, ED-54, pm) Number of sectors: 1.5 allia size: 0.800 gFlua) Explore ET-Dusty	CMFGEN Besed on CMFCD4 model atmospheres UV, WS and intranges	CMFCEN-OB-LMC-24 Eved of CMFCEN robot is annupper UV 9009 5 30 000 AV01 900 000 151 000 Av1 and ((1 000 162 15 000 Av1) spectral togost frages
CMFGEN-08-LMC-24-SED Bacd on CMFCDH model almospheres PMFC [Insert Almospheres] PMFC[Insert Almosphere] [Insert [Insert Almosphere] PMFCB [Insert Almosphere] [Insert Almosphere] Penderode by JC. Borrel (LAM Marsile) Number of spectra [Insert Alm Marsile) Laplace CMFCDH-08-LMC-24-8ED Download	CMFCEN-08-SMC-24 Berd on CMFCR model atmospheri UV 9000 to 300 AVV 90 00 the 120 AVA and RR (12 000 to 25 000 AVA) spectral ranges Typiff: Bioric Knond [I dage E Biose Knone] Provide (J bioric Knond [I dage E Biose Knone] Provide (J bioric Knond [I dage E Biose Knone] Provide (J bioric Knond [I dage Biose Knone] Provide (J bioric Knond [I dage Biose Knone] Provide (J bioric Knond [I dage Biose Knone] Explore (J biose Knone] Explore (J biose Knond [I dage Biose Knone] Explore (J biose Knone] Explore (CMEGEN-DB-SMC-24-SED Based on CMEGEN model atmospheres TyPE The Smarth S	CMFGEN-SED Based on CMFGEN model atmospheres MyRef (22224): 52377(1) (adge (2.234, 4.18)) (%)(4.16, 1.26)) (br = 15 km) Probaced by Martina (LUPM, Montpetiin) Number of adjects (1.26 kuise (2.6440 (FM)) (Laguere CMFGEN 42D) Desminat
CMFCEN-WR Bard on OVFCDH model Attransplares V5 ranges P* (5) 754, 754000 (1 sg /Law) (5 /S. 54) Pendice (1 /S. 40400 (1 sg /Law) (5 /S. 54) Pendice (1 /S. 54)	CMFCEN-WR-SED Basel on OVFCRN model Atmospheres VS ranges P* C [140,0.00] (arg (Z.taw) 6 [3.3.5.3] Pyr01 (b (0.0.0) 0* T 51m) 0* T 51m) Nonbert of XD 2] [140 (are: 0.048 (Hal) Databert of XD 2] [140 (are: 0.048 (Hal) Databert of XD 2] [140 (are: 0.048 (Hal)	PLATO RESTRUCTED ACCESS Based on MARCS model exmoghners and TURBOSPECTRUM NUTE 2.8 spectral updatesis VIS mores Republic Plane 2010 Provide Tax a 2010 Resolution - 2000 Produced by the PLATO collaboration NP 122100 Number of spectra - 2 Toxis are 0 MMB Plane	PLATO-INTENSITY ESSINCTED ACCESS Intensity spectra based on MARCS model atmospheres and TURBOSPECTRUM NOTE 2.0 spectra synthesis fulfy (= (4533, 45116) (Large (=), 4, 151) /Pulf (= (4533, 45116) (Large (=), 4, 151) /Pulf (= (4532, 45116)
PLATO-SED	POPSYCLE	RSG	STAGGER

Figure 12: POLLUX Collection Query Interface.

From the collection page, the user may either retrieve the entire collection as an archive file (the size of which is given as the total size of the collection), or explore the collection.

In the second case, the user will access the stellar parameter query interface (described hereafter) that is also directly accessible from the homepage, with a pre-selected query tree for the collection that was chosen.

As of the current 12th release, private collections also exist (PLATO, PLATO-INTENSITY, PLATO-SED). For unauthorized visitors, caracteristics of the collection are visible but the collection cannot be queried nor explored. The access is granted upon request and validation by the consortium proprietary of the original data. Once authorized, the users have to log in to unlock the collections.

4.2 Stellar Parameters Query Interface

4.2.1 Query of synthetic spectra

This query is hierarchical and the parameters that can be queried in the right part of the page are adjusted according to the choice of spectra type, collection, spectral domain, and model type made in the left part of the query page.

The default display depends on the access path to this page. If coming from the homepage of POL-LUX, all synthetic spectra are selected for query except the spectra of the private collections and the WR spectra for which there is a special query interface, as shown in Fig. 13. If coming from a collection, the General parameters are already preselected to limit the query to this selected (see Fig 14).

The choice of the **General Parameters** on the left is made by clicking on the relevant boxes. The selected boxes appear in green. The selections are automatically exclusive : for instance, if the

THE SYNTHETIC STELLAR SPECTRA DATABASE POLLUX	SPECTRA COLLE	CTIONS STE	LLAR PARAMETERS	USER GUIDE	S↓ MY SPEC	CTRA [→
EX	PLORE BY STELLA	R PARA	METERS			
General parameters:	Spectra variables:					
Spectra type: Synthetic Spectra SED	Spectrum parameters	Lowest	Lower	Equal	Upper	Highest
Collection: AMBRE BT-Dusty CMFGEN	Effective temperature : K	2000				63880
	Gravity log10: cgs	-0.5				5.5
CMFGEN-OB-LMC-24 CMFGEN-OB-SMC-24	Microturbulent velocity ξ_t : km/s	0				10
CMFGEN-WR PLATO	Metallicity : [Fe/H]	-5				1
PLATO-INTENSITY POPSYCLE RSG	Specific Abundances					
STAGGER STAGGER-INTENSITY	Carbon : [C/Fe]	-1.31				0.03
STAGGER-RVS	Nitrogen : [N/Fe]	-0.01				1.2
Spectral domain: UV VIS IR						
Model type: 1-D Plane Parallel (p) Spherical (s)						
3-D 3D RHD						
Search						
	ERSITÉ DE ORE	Em	CNTS		GS	

Figure 13: POLLUX Stellar Parameter Specific Query Interface.

AMBREcollection is the only one selected, the boxesSynthetic SpectraVISPlane-ParallelandSphericalwill be automatically selected while all the others remain white (can still be selected) or grey (incompatible with the currently selected collection) as illustrated in Fig 14.

The user can specify the **Spectra Variables**. The possible query depends on the selected **General Parameters** that has been chosen :

- effective temperature
- log g
- mass (appears only when spherical models are selected)
- luminosity (appears only when spherical models are selected)
- micro-turbulent velocity
- metallicity [Fe/H]

When selecting the **CMFGEN-WR** collection, a specific query form will appear (as shown in Fig. 15) in which the parameters that can be queried are different :

- temperature at optical depth $\tau = 20 \text{ T}^*$
- luminosity
- micro-turbulent velocity
- hydrogen mass fraction X

THE SYNTHETIC STELLAR SPECTRA DATABASE POLLUX	SPECTRA COLLECTIONS	STELLAR PARAMETERS	USER GUIDES ↓	MY SPECTRA
EXP	LORE BY STELLAR PA	RAMETERS		
General parameters: 🕕	Spectra variables:			
Spectra type: Synthetic Spectra SED	Spectrum parameters Low	est Lower	Equal Upper	Highest
Collection: AMBRE BT-Dusty CMFGEN CMFGEN-OB-LMC-24 CMFGEN-OB-SMC-24 CMFGEN-WR PLATO PLATO-INTENSITY POPSYCLE RSG STAGGER STAGGER-INTENSITY STAGGER-RVS	Effective temperature : K 200 Gravity log_{10} : cgs 0. Microturbulent velocity $\xi_i: kn/s$ 0.11 Metallicity: $[Fe/h^{j}]$ 0 Specific Abundances 0 Alpha elements: $[a/Fe]$ 0 Carbon: $[C/Fe]$ 0 Oxygen: $[a/Fe]$ 0			6000 5.5 1.139 0 0 0 0
Model type: 1-D Plane Parallel (p) Spherical (s) 3-D 3D RHD Search	Nitrogen: ///Fe/ 0	crrs	G	

Figure 14: Example of exclusive query.

EXF	PLORE BY STELLA	R PARA	METERS			
General parameters: 🚯	Spectra variables:					
pectra type: Synthetic Spectra SED	Spectrum parameters	Lowest	Lower	Equal	Upper	Highest
llection: AMBRE BT-Dusty CMFGEN	Temperature@Tau=20 T* : K	33780				74300
	Mass : solar mass	10				30
CMFGEN-OB-LMC-24 CMFGEN-OB-SMC-24	Luminosity log ₁₀ : L _{sun}	5.3				5.9
CMFGEN-WR PLATO	Microturbulent velocity ξ_t : km/s	10				30
PLATO-INTENSITY POPSYCLE RSG	X : Hydrogen mass fraction	0				0.5541
	Y : Helium mass fraction	0.3827				0.9923
STAGGER STAGGER-INTENSITY	Metallicity : [Fe/H]	0				0
STAGGER-RVS	Specific Abundances					
pectral domain: UV VIS IR	Alpha elements : [a/Fe]	0				0
odel type: 1-D Plane Parallel (p) Spherical (s)	Carbon : [C/Fe]	-2.39				3.21
	Oxygen : [O/Fe]	-2.66				2.34
3-D 3D RHD	Nitrogen : [N/Fe]	0				1.92
Search	X _C : Carbon mass fraction	2.93e-6				0.459
Staten	X ₀ : Oxygen mass fraction	3.91e-6				0.153
	X _N : Nitrogen mass fraction	0				0.0171

Figure 15: Query form for spectra in the CMFGEN-WR collection

- helium mass fraction Y
- metallicity [Fe/H]

The user can either choose an interval or choose an exact value for the **Spectra Parameters**. The extremum values for each parameter in the selected spectra are also given.

A second query block is also available, which enables the user to choose data sets with **Specific Abundances**. The set of searchable abundances depends on the data queried. For CMFGEN and STAGGER data, queries can be made on Carbon, Nitrogen, Oxygen and α -elements in terms of [X/Fe]. For the AMBRE data, *r*- and *s*- elements abundances in terms of [X/Fe] can also be queried. For the POPSYCLE data, the mass fraction of He Y can be queried. For CMFGEN-WR data, the searchable abundances are Carbon, Nitrogen and Oxygen in terms of mass fractions X_C, X_N and X_O.

Once the choices made, the request is sent by clicking on the Search button below the list of **General Parameters** on the left.

- If the exact value requested is not available in the database, a "no spectra" message will be returned.
- If the values queried exist, the spectra complying with the query will be displayed in the lower part of the window.
- If no value is entered in the **Spectra Parameters** fields, all the models complying with the selected **General parameters** will be displayed in the lower part of the window.

4.2.2 Query of spectral energy distributions

The query of form fro SEDs, currently only available for CMFGEN data in the public database, is similar to that of synthetic spectra as shown in Fig. 16.

THE SYNTHETIC STELLAR SPECTRA DATABASE POLLUX	SPECTRA COLLECTIONS	STELLAR PARAMETERS	S USER GUIDES	S↓ MY SPECTRA [C→
EXF	PLORE BY STELLAR PA	RAMETERS			
General parameters: 🕕	Spectra variables:				
Spectra type: Synthetic Spectra SED	Spectrum parameters Low	vest Lower	Equal	Upper Highest	
Collection: CMFGEN-OB-LMC-24-SED	Effective temperature : K 25	00		63880	
CMEGEN-OB-SMC-24-SED CMEGEN-SED	Gravity log ₁₀ : cgs 2	.4		5.5	
	Mass : solar mass	0		403.81	
CMFGEN-WR-SED PLATO-SED	Metallicity : [Fe/H] -1.	48		1	
Model type: 1-D Plane Parallel (p) Spherical (s)	Specific Abundances				
3-D 3D RHD					
Search					
	RSITÉ DE ORF	Cnrs			
	PELLIER			DESERVATORE VIRTUEL CRAND SUD QUEST	

Figure 16: Query form for SEDs.

4.3 Results of Request

The result of the request is displayed on the lower part of the query page and consists of a table (dispatched on several pages if needed) containing 4 main parts : **Cart**, **Display**, **Download**

and **Data Characteristics** as illustrated in Figure 17.

The table may be sorted according to a specific field by clicking on the column name.

There	are 13	89 synth	etic spectra (SPEC) co	orresponding to th	nese paramete	rs:	Add	all results to "M	IY SPECTR	A"		Remove al	ll results from	MY SPECTR	۹"	Display	20 v spect	ra per page
Ô		⊡	Collection	Model	Range	Туре	Specs 🔒	Teff	Logg	ξt	Y	[Fe/H]	[a/Fe]	[C/Fe]	[O/Fe]	[N/Fe]	[r/Fe]	[s/Fe]
Ä		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10000	3.5	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10000	3.5	1	0.256	-0.77	0.4	0	0.4	0	0	0
Ä		÷	POPSYCLE	ATLAS12	VIS	P	91900002	10000	4	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10000	4	1	0.256	-0.77	0.4	0	0.4	0	0	0
Ä		÷	POPSYCLE	ATLAS12	VIS	P	91900002	10000	4.5	1	0.256	-0.77	0.4	0	0.4	0	0	0
嶌		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10000	4.5	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
Ä		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10250	3.5	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10250	3.5	1	0.256	-0.77	0.4	0	0.4	0	0	0
Ë		Ł	POPSYCLE	ATLAS12	VIS	P	91900002	10250	4	1	0.256	-0.77	0.4	0	0.4	0	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10250	4	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	P	91900002	10250	4.5	1	0.256	-0.77	0.4	0	0.4	0	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	P	91900002	10250	4.5	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10500	3.5	1	0.256	-0.77	0.4	0	0.4	0	0	0
E		Ŧ	POPSYCLE	ATLAS12	VIS	P	91900002	10500	3.5	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
Ä		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10500	4	1	0.256	-0.77	0.4	0	0.4	0	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10500	4	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
ä		Ŧ	POPSYCLE	ATLAS12	VIS	P	91900002	10500	4.5	1	0.256	-0.77	0.4	0	0.4	0	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	Р	91900002	10500	4.5	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	P	91900002	10750	4	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
Ë		Ŧ	POPSYCLE	ATLAS12	VIS	P	91900002	11250	3.5	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
_	_	_					<	< Page	1/70	>	>							

Figure 17: POLLUX Result of Request for synthetic spectra. The first column allows to upload all or selected spectra into the "MY SPECTRA" space; the second column allows to display the spectrum and its header; the third column allows to directly download the selected spectra; the remaining columns give the attributes of the spectra retrieved in terms of General and spectra parameters.

The number of synthetic spectra found and the number of pages are also indicated. The User may also change the number of spectra displayed per page, add or remove all the results to the "MY SPECTRA" workspace that is also accessible via the top menu of the query page.

4.3.1 Cart, Display and Download

The three first columns of the query result correspond to the cart, the display and download of the spectra:

1. ① The first column of the query result allows to store the selected spectra into the workspace from which further visualization, process and download is possible (see below). When clicking on the empty cart icon 몇, the icon will change to 몇 to indicate that the selected item has been uploaded to the "MY SPECTRA" workspace ⁵. On the top menu of the page, the "MY SPECTRA" section will indicate the number of spectra that are available in the workspace, as illustrated in Fig. 18.

⁵This operation can be reverted by clicking again on the 🙀 icon which will change to the 🙀 , and remove the spectrum from the MY SPECTRA workspace.

THE SYNTHETIC STELLAR SPECTRA DATABASE POLLUX						SPECTRA	A COLLECTION	5 STEL	LAR PARAMETER	S USER GU	IDES ↓	MY SPECTR	A [→
	E	EXPLORE BY STELLAR PARAMETERS											
General parameters:		Spectra variables:											
Spectra type: Synthetic Spectra SED		Spectrum parameters			Lo	owest	Lowe	r	Equal	Up	er	Highe	st
Collection: AMBRE BT-Dusty CMFGEN CMFGEN-OB-LMC-24	Ef	fective temperature : K			3	8800						1400	0
	Gr	avity log10: cgs				1						5.5	
CMFGEN-OB-SMC-24 CMFGEN-WR PLATO	мі	croturbulent velocity &:	km/s			1						1	
PLATO-INTENSITY POPSYCLE RSG STAGGER	Y	Y : Helium mass fraction			0	.256						0.3	
STAGGER-INTENSITY STAGGER-RVS	M	Metallicity : [Fe/H]			-	1.58						-0.4	,
Spectral domain: UV VIS IR		Specific Abundances)										
	AL	Alpha elements : [a/Fe]				0.4						0.4	
Model type: 1-D Plane Parallel (p) Spherical (s)	Ca	Carbon : [C/Fe]			-0.3							0	
3-D 3D RHD	0)	ygen : [O/Fe]			-0.05							0.4	
	Ni	trogen : [N/Fe]				0						1.2	
Search	R	process elements : [r/Fe]				0						0	
	Sp	process elements : [s/Fe]				0						0	
There are 1389 synthetic spectra (SPEC) corresponding to these parameters :		Add all res	ults to "MY SP	ECTRA*			Remove all r	esults from	"MY SPECTRA"		Disp	olay 20 - spectr	a per page
🖞 🕩 🕢 Collection Model Range	Туре	Specs 🔒	Teff	Logg	ξt	Y	[Fe/H]	[a/Fe]	[C/Fe]	[O/Fe]	N/Fe]	[r/Fe]	[s/Fe]
POPSYCLE ATLAS12 VIS	р	91900002	10000	3.5	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
📔 🖬 📩 POPSYCLE ATLAS12 VIS	р	91900002	10000	3.5	1	0.256	-0.77	0.4	0	0.4	0	0	0
POPSYCLE ATLAS12 VIS	P	91900002	10000	4	1	0.3	-0.77	0.4	-0.3	-0.05	1.2	0	0
ELI . POPSYCLE ATLAS12 VIS	P	91900002	10000	4	1	0.256	-0.77	0.4	0	0.4	0	0	0

Figure 18: Example selected spectra uploaded in the "MY SPECTRA" workspace as seen from the query page.

2. D The second column of the query result provides access to a new window by clicking on thea icon, that opens as a separate tab in the web browser, where the spectrum is displayed alongside its header.

The plotting interface is shown in Fig 19. The name of the spectrum is recalled as a global title of the page.

It contains two parts: the spectrum plot on the left and the header on the right. Both the absolute flux and normalized flux are displayed as a function of wavelength in Å in two distinct clickable tabs (the background of the selected is of the same colour as the spectrum on the plot). In the case of SEDs, there is only one tab, the background of which is also of the same colour as the spectrum on the plot.

The plots are interactive : the User may choose a logarithmic display for the ordinates (Change y scale), enter the wavelength range of the chart below the plot, or simply zoom with the mouse directly on the plot.

The plot can be reset to default boundaries at any time by clicking on the Reset zoom button. It can also be enlarged (and reduced) by clicking on the C icon.

The displayed plot can also be saved as a .png file using the Save to PNG button.

From this display page, the spectrum can be either downloaded Download or added to the "MY SPECTRA" workspace Add to my spectra, with the buttons right above the header file display.



Figure 19: Example of spectrum and header display.

3. The third column of the query result allows the direct download of the spectrum in various formats. Clicking on the 🕹, a pop-up appears as in Fig. 20, where the pre-selected download format is Flat Table in zip format, and other formats can be also chosen (FITS, VoTable, XML Table, XML Binary VoTable) before actually downloading.

Colle	ction:	AMB	RE BT-Dusty	CMFGEN	E	ffective tempe	rature : K	2	000						6000
					G	ravity log ₁₀ : cg	7 5		0.5						5.5
		CMF	GEN-OB-LMC-24	CMFGEN-OB-SMC-	N	licroturbulent	velocity ξ _t : km/s	0	.104						1.139
		CMF	SEN-WR PLATO		M	etallicity : [Fe/	′H]		0						0
		PLAT	O-INTENSITY PC	OPSYCLE RSG		Specific Abu	ndances 🕕								
		STAC	GER STAGGER-IN	NTENSITY	A	lpha elements	: [a/Fe]		0						0
		STAG	GER-RVS		с	arbon : [C/Fe]			0						0
					c	xygen : [O/Fe]			0						0
Spect	ral dor	main:	UV VIS IR		N	itrogen : <i>[N/Fe</i>	1		0						0
							-								
Mode	a type:	1-D	Plane Parallel (p)	Spheric		DOWN	LOAD THI	S SPE	CTRU	M	\times				
		3-D	3D RHD												
						elst estels					- The last				
			Search	Dat	a rormac:	Flat Table	FIIS			ML Binary v	orable				
				D	ownload										
There	e are 75	54 synth	etic spectra (SPEC) co	orrespondi											
Ê	▶			hicspondi								'MY SPECTRA	Displa	ay 20 ~ spec	ra per page
Ä		٩	Collection	Model	Range	Туре	Specs 🚯	Teff	Logg	ξt	[Fe/H]	'MY SPECTRA" [a/Fe]	Displa	ay 20 v spect	ra per page [N/Fe]
	-	 ④ 	Collection BT-Dusty	Model PHOENIX	Range VIS	Type s	Specs () 91101091	Teff 4700	Logg 4.5	ξ t 0.479	[Fe/H] 0	MY SPECTRA" [a/Fe] 0	Displa	ay 20 v spect [O/Fe] 0	ra per page [N/Fe] 0
Ä	4	(← [←	Collection BT-Dusty BT-Dusty	Model PHOENIX PHOENIX	Range VIS VIS	Type s s	Specs () 91101091 91101091	Teff 4700 4800	Logg 4.5 5	ξ t 0.479 0.493	[Fe/H] 0 0	'MY SPECTRA" [a/Fe] 0 0	Displa [C/Fe] 0 0	ay 20 - spect [O/Fe] 0 0	ra per page [N/Fe] 0 0
iă iă	1	(← [← [←	Collection BT-Dusty BT-Dusty BT-Dusty	Model PHOENIX PHOENIX PHOENIX	Range VIS VIS VIS	Type S S S	Specs ① 91101091 91101091 91101091	Teff 4700 4800 4900	Logg 4.5 5 0.5	ξ t 0.479 0.493 0.528	[Fe/H] 0 0	'MY SPECTRA" [a/Fe] 0 0 0	Displa [C/Fe] 0 0 0	ay 20 ~ spect [O/Fe] 0 0 0	ra per page [N/Fe] 0 0 0
יע יע יע	11	(+ [+ [+ [+	Collection BT-Dusty BT-Dusty BT-Dusty BT-Dusty	Model PHOENIX PHOENIX PHOENIX PHOENIX	Range VIS VIS VIS VIS	Type S S S S	Specs 91101091 91101091 91101091 91101091 91101091 91101091	Teff 4700 4800 4900 4900	Logg 4.5 5 0.5 1	ξ 0.479 0.493 0.528 0.528	[Fe/H] 0 0 0	MY SPECTRA"	Displa [C/Fe] 0 0 0 0	ay 20 ∨ spect [O/Fe] 0 0 0 0	[N/Fe] 0 0 0 0 0 0
ात :ात :ात		(+ [+ [+ [+	Collection BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty	Model PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX	Range VIS VIS VIS VIS VIS VIS	Type S S S S S S	Specs Image: The system 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091	Teff 4700 4800 4900 4900 4900	Logg 4.5 5 0.5 1 1.5	ξ 0.479 0.493 0.528 0.528 0.528	[Fe/H] 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	MY SPECTRA" [a/Fe] 0 0 0 0 0 0 0	Displa [C/Fe] 0 0 0 0 0 0 0 0 0	ay 20 ✓ spect [O/Fe] 0 0 0 0 0 0	[N/Fe] 0 0 0 0 0 0 0 0
ים ים ים ים		(د (د (د (د	Collection BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty	Model PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX	Range VIS VIS VIS VIS VIS VIS VIS	Type S S S S S S S	Specs (1) 91101091 91101091 91101091 91101091 91101091 91101091	Teff 4700 4800 4900 4900 4900 4900 4900	Logg 4.5 5 0.5 1 1.5 2	ξ t 0.479 0.493 0.528 0.528 0.528 0.528	[Fe/H] 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	MY SPECTRA" [a/Fe] 0 0 0 0 0 0 0	Displa [C/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	ay 20 ∨ spect [0/Fe] 0 0 0 0 0 0 0 0 0 0 0	ra per page [N/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0
יע ייע ייע ייע ייע		(+ (+ (+ (+ (+)	Collection BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty	Model PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX	Range VIS VIS VIS VIS VIS VIS VIS VIS	Type S S S S S S S S S S	Specs Image: Constraint of the system 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091	Teff 4700 4800 4900 4900 4900 4900 4900 4900 49	Logg 4.5 5 0.5 1 1.5 2 3	ξ _t 0.479 0.493 0.528 0.528 0.528 0.528 0.528 0.528 0.528	[Fe/H] 0 0 0 0 0 0 0 0 0	MY SPECTRA" [a/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	Displa [C/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	ay 20 ∨ spect [0/Fe] 0 0 0 0 0 0 0 0 0 0 0	(N/Fe) 0 0 0 0 0 0 0 0 0 0 0 0
ימ ימ ימ ימ ימ ימ		(د (د (د (د (د (د (Collection BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty	Model PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX	Range VIS VIS VIS VIS VIS VIS VIS VIS	Type S S S S S S S S S S S S	Specs Image: Control of the system 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091	Teff 4700 4800 4900 4900 4900 4900 4900 4900 49	Logg 4.5 5 0.5 1 1.5 2 3 4	ξ 0.479 0.493 0.528 0.528 0.528 0.528 0.528 0.528	[Fe/H] 0 0 0 0 0 0 0 0 0 0 0 0 0	MY SPECTRA" [a/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	Displa [C/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	ay 20 ∨ spect [0/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	ra per page [N/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0
ימ ימ ימ ימ ימ ימ ימ		لد (د (د (د (د (د (د و	Collection BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty	Model PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX PHOENIX	Range VIS VIS VIS VIS VIS VIS VIS VIS VIS	Type S S S S S S S S S S S S S S S S S	Specs Image: Control of the system 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091	Teff 4700 4800 4900 4900 4900 4900 4900 4900 49	Logg 4.5 5 0.5 1 1.5 2 3 4 4.5	ξ _t 0.479 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528	[Fe/H] 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	MY SPECTRA" [a/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	Displa 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	ay 20 ∨ spect [O/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	rra per page [N/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0
ימ ימ ימ ימ ימ ימ ימ ימ		لد لد لد لد لد لد له له ف	Collection BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty BT-Dusty	Model PHOENIX	Range VIS VIS VIS VIS VIS VIS VIS VIS VIS VIS	Type S	Specs 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091 91101091	Teff 4700 4800 4900 4900 4900 4900 4900 4900 49	Logg 4.5 5 0.5 1 1.5 2 3 4 4.5 0.5	ξε 0.479 0.493 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528 0.528	[Fe/H] 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	MY SPECTRA" (a/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	2 Displ 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	ay 20 ∨ spect [O/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0	ra per page [N/Fe] 0 0 0 0 0 0 0 0 0 0 0 0 0

Figure 20: Direct download of a spectrum

In order to clearly acknowledge the producers of the data (HRSS or SEDs), a dedicated README.txt file for each collection (an example is given hereafter) is systematically added to the archive to be downloaded in which explicit names and papers to be cited along with the POLLUX database are given.

You will find here the synthetic stellar spectra you have selected and downloaded from the POLLUX database (pollux.oreme.org). They were generated by P. De Laverny (OCA) within published scientific work and is to be acknowledged accordingly : AMBRE : de Laverny, P.; Recio-Blanco, A.; Worley , C.C. & Plez, B., 2012, A& A 544, A126

The database can be also acknowledged with Palacios et al., 2010, A& A 516, A13.

You will find all relevant information concerning the data, the codes and the inputs used to generate them, as well as detailed curation information in the header file attached to each spectrum.

Thank you for using the service!

The POLLUX DB Team.

4.3.2 Spectrum Parameters

Columns 4 to 21 of the table present the data characteristics. It adapts to the selected General parameters and can differ from one collection to another. When the query concerns several collections, only the common data characteristics to all the selected collections are displayed. In any case, columns 4 to 8 are always the following :

- 4. Collection Subset to which the spectrum belongs
- 5. Model Radiative transfer code used to compute the spectrum
- 6. Range Spectral Domain
- 7. Type spherical, plane-parallel or 3D RHD model atmosphere
- 8. Specs Specifications of the synthetic spectrum as defined in Tab. 4.

The remaining columns will depend on the collections selected and can be

- 9. T_{eff} Effective temperature in K
- 10. log g Surface gravity in log
- 11. Mass Stellar mass in M_{\odot} units only relevant for spherical models
- 12. Lum Logarithm of the stellar luminosity expressed in L_{\odot} units only relevant for spherical models
- 13. ξ_t Micro-turbulent velocity in km.s⁻¹
- 14. [Fe/H] Metallicity with respect to solar
- 15. **[C/H]** Carbon abundance
- 16. [N/H] Nitrogen abundance
- 17. **[O/H]** Oxygen abundance
- 18. $[\alpha/H] \alpha$ elements abundance
- 19. [s-elements/H] abundance of s-elements
- 20. [*r*-elements/H] abundance of *r*-elements
- 21. Y helium mass fraction

Clicking on the label of these columns will sort the data in the table in decreasing or increasing values of the selected parameter.

For the specific case of Wolf-Rayet spectra (CMFGEN-WR collection); the data characteristics of columns 9 to 18 are the following

- 9. **T**^{*} temperature at optical depth $\tau = 20$
- 10. Lum Logarithm of the stellar luminosity expressed in L_{\odot} units
- 11. ξ_t Micro-turbulent velocity in km.s⁻¹
- 12. X hydrogen mass fraction
- 13. Y helium mass fraction
- 14. [Fe/H] Metallicity with respect to solar
- 15. X_C carbon mass fraction
- 16. X_N nitrogen mass fraction
- 17. X_O oxygen mass fraction

4.4 The "MY SPECTRA" workspace

Newly introduced in the present version of the database portal, the workspace is where data analysis and download is meant to be done. It is presented in Fig. 21.

Once in this area, you can return to the selection form and results of request by using the back button of your browser.

Pal	THE SYNTHETIC STELLAR SPECTRA DATABASE POLLUX	SPI	ECTRA COLLECT	TIONS	STELLAR PARAMET	ERS USER'S GU	IIDE MY SPECTRA		
		MY SPECT	FRA						
	You saved 4 spectra and 0 convolved spectra :								
	Download Overplot Conve	olution Provenance	Compan	ion Dat	a SAMP-VO	Remove			
- 6	Name	Author	Date	Туре	Collection	Spectral domain	Atmosphere model		
0	M_s4500g0.5z0.00t1.0_a0.00c0.00n0.00o0.00r0.00s0.00_IR.spec	P Hernandez Cascales	Feb. 5, 2022	SPEC	RSG	IR	MARCS		
	M_p4500g5.0z0.00t1.0_a0.00c0.00n0.00o0.00r0.00s0.00_VIS.spec	de Laverny	Feb. 1, 2010	SPEC	AMBRE	VIS	MARCS		
	P_s3900g3.50z0.0t0.37_a0.00c0.00n0.00o0.00_IR.spec	F. Allard	Sept. 13, 2017	SPEC	BT-Dusty	IR	PHOENIX		
	O_3d3899g2.00z-0.50_a0.20c0.00n0.00o0.20_mu0.90_UV.spec	A. Chiavassa	Jan. 1, 2017	SPEC	STAGGER-INTENSITY	UV	STAGGER		
		ORE	ĔŇ		cnrs	DESERVATORE			

Figure 21: The MY SPECTRA workspace

The names of the spectra in the "MY SPECTRA" page can be clicked : this will open a new tab in the web browser containing the plot and the display of the spectrum.

Moreover, four functionalities are available from the workspace :

1. Download

From the MY SPECTRA workspace, the selected spectra can be downloaded (default download format will be Flat table and FITS, VoTable, XML, XML VoTable and XML Binary VoTable are also available for selection) with the same pop-up interface as from the result of query page described illustrated in Fig. 20.

2. Overplot



STELLAR SPECTRA OVERPLOT



Figure 22: Overplot of three spectra from the MY SPECTRA workspace

The selected spectra in the workspace can be overplotted. The plot will appear in a new tab of the web browser. Each spectrum on the overplot is assigned a colour, and the different tabs on the right part of the window are coloured accordingly. The header of the corresponding spectrum is displayed when clicking on the corresponding colour tab. Per default, the displayed header is the one of the first spectrum as listed in the legend of the plot (Spectrum 1). Moving the mouse over the corresponding coloured rectangle in the legend will display the name of the spectrum in a colour framed box above the plot space.

The plot functionalities are as described before. There is an additional possibility of changing the order of the plotted spectra to improve clarity by clicking on the colour of the spectrum in the Bring to Front section below the plot. Clicking directly on the name of the spectrum in the legend of the plot will moreover hide the selected spectrum. It will be shown again by clicking a second time on its name.

3. Convolution

The interface provided by clicking on this button, and shown in Fig. 23 allows the query of the SPECONVOL VO service based on a Fortran code that implements a number of convolution functions used to model the instrumental profile, the macroturbulence and the rotational broadening of spectral lines.

THE SYNTHETIC STELLAR SPECTRA DATABASE POLLUX				SPECTRA COLLECTION	S STELLAR PARAMETERS	USER'S GUIDE	MY SPECTRA
	CON	VOLU	JTION SEF	RVICE			
	You	have se	elected 1 spect	:rum:			
	Name M_p6000g5.0z0.50t1.0_a-0.40c0.00n0.00)o-0.40r0.00s0.	.00_VIS.spec				
	Macroturbulence velocity :	0.0	Unit: km/s ~				
	Rotational velocity :	0.0	Unit: km/s ~				
	Instrument profile :	0.0	Unit: mA 🗸				
	Radial velocity :	0.0	Unit: km/s ~				
	Central wavelength :	8525	Unit: A 🗸				
	Width :	Apply	Unit: A 🗸				
UNIVER'S & REALECTERS	UNIVERSITÉ DE MONTPELLIER		ORE~3	leans o	cors		O BOL

Figure 23: Interface to the SPECONVOL convolution service.

The user can choose to include one to four of the following convolution parameters:

- Macroturbulence velocity can be specified in km.s⁻¹. It is modelled by a radial tangential anisotropic profile as described in Gray (2005, pp 433), and describes the lines broadening due to convection;
- Rotational velocity of the star can be specified in km.s⁻¹. It is modelled by a "rotation" profile as described in Gray (2005, pp 434-436), and describes the lines broadening due stellar rotation;
- The simulated signature of an instrumental profile can be specified in mÅ (default unit) or in km.s⁻¹. It is modelled by a Gaussian profile and describes the line broadening due to the instrumental design.

The convolution parameters are applied to a spectral window centred at a chosen central wavelength that can be specified in units Å, and of limited width to be chosen between 100 Å and 500 Å.

The convolved portion of spectra can also be translated according to the a radial velocity that can be specified in units of km.s⁻¹.

These parameters are sent to the SPECONVOL service when clicking on the apply button and the user is redirected after a short time to the workspace page that now includes the initially selected spectra and the associated convolved portions.

A new name is given to the convolved spectrum, integrating the parameters of convolution as illustrated in Fig. 24



Figure 24: Example of nomenclature for a convolved portion of a spectrum using SPECONVOL.

The original spectrum and the convolved portion can be overplotted using the graphical tool within the interface as illustrated in Fig. 25.



Figure 25: Rendering of overplotted spectrum and convolved portion.

4. Provenance

The interface provided by clicking on this button, and shown in Fig. 26 allows to obtain the Provenance of the spectrum following the ProvenanceDM and ProvSAP protocol of the IVOA. The provenance refers to the census of all the information related to the entities, activities and authors involved in the collection, creation, modification or sharing of data. The interface provided allows to specify the parameters for the ProvSAP protocol as of 2018, which is a working draft. The result of query is either a raw output (in JSON, XML) or a graphical

output (in SVG, PNG or PDF). The direction of the provenance can only be backwards. The "agent" parameter includes information concerning the authors of the data. The "model" parameter corresponds to the Provenance model used (either from IVOA⁶ or W3C⁷). Finally, the meaning of "depth" parameter is illustrated in Fig 27.

THE SYNTHETIC STELLAR SPECTRA DATABASE POLLUX	SPECTRA COLLECTIONS	STELLAR PARAMETERS	USER GUIDES ↓	MY SPECTRA	[•
	PROVENANCE	SERVICE			
M_s5000g1.0z-0.25t2.0_a0.30c0	Spectrum key .00n0.00o0.30r0.00s0.00_conva1.5cc	nvny8.0convy0.076convb0.5ml	0.0mb0.0_VIS_V10.spec		
	Depth : Resource Format :	1 v			
	Direction :	BACK ~			
	Model:	IVOA ~			
	Apply				
			rs 🖸		



Figure 27: Definition of the DEPTH parameter for ProvSAP

Figure 26: Interface to the Provenance Visualisation service

5. Companion Data

This button gives access, for some data only (as of the current release, companion data exist for the BT-Dusty, AMBRE, RSG, POPSYCLE public collections). The companion data are either (i) atomic or molecular linelists that are associated to the spectrum (used in the workflow to compute it) (ii) model atmospheres used in the workflow to generate the spectra. Clicking on this button will open a menu with one or both of these resources and either directly give access to the companion file, or open the database hosting the companion data.

6. SAMP-VO

This button is unlocked for one spectrum at the time. It allows to broadcast the selected spectrum to a VO-compliant application with a SAMP hub such as Cassis⁸ or TOPCAT⁹. To use it, the SAMP hub must be opened. Clicking on the button will automatically broadcast the spectrum to the application, say Cassis, that will in turn open a window indicating that the data was received and allowing the user to visualize/modify/treat it within that application.

7. Remove

This last button allows to remove spectra from the workspace.

⁶https://ivoa.net/documents/ProvenanceDM/

⁷https://www.w3.org/TR/prov-dm/

⁸cassis.irap.omp.eu

⁹https://www.star.bris.ac.uk/~mbt/topcat/

5 FUTURE DEVELOPMENTS

5.1 ATLAS data to fill the gap of warm stellar spectra

The archive will be completed with corrected data sets derived from ATLAS model atmospheres at various metallicities.

5.2 Extension of the database to IR

Considering the new generation of spectrographs that are being built, a new effort to provide theoretical data in the IR domain should be done and we are working on this aspect in order to be able to provide the community with well described ans VO-compliant IR high resolution synthetic spectra.

5.3 (Re-)Introduction of SEDs

The introduction of spectral energy distributions (SED data) for warm and cool stars will be considered in the future.

5.4 Companion data and associated services

The companion data of our spectra are, in some cases, distributed in other databases that are not VO-compliant at present. It is the case of the MARCS model atmospheres database (marcs.oreme. org) for instance. In the mid-term future we plan to actually add an interoperability layer to be able to directly retrieve the exact companion model atmosphere to a spectrum in the POLLUX database instead of just pointing to the other service.

5.5 Development of an interface to query and run MARCS/Turbospectrumv2.0 from the POLLUX web interface

In a longer term, and in order to provide maintained access to the codes MARCS and Turbospectrum on a long term, we plan to develop an interface to run these codes from the webpage on demand. This will allow users to have tailored made models and to feed the database with their computation, as is it done in other services.

6 CREDITS

The POLLUX database is described at length as of its third release, in a Palacios et al. (2010). When using POLLUX data for scientific publication, please quote :

Palacios A., Gebran M., Josselin E., Martins F., Plez B., Belmas M., Lèbre A., 2010, A&A 516, A13

and mention the following sentence:

This research was achieved using the POLLUX database (http://pollux.oreme.org/), operated at LUPM (Université de Montpellier - CNRS, France) with the support of the PNPS and INSU.

7 **REFERENCES**

- 1. Allard, F., Homeier, D., Freytag, B., Sharp, C. M., 2012, EAS Publications Series, 57 3-43 "Atmospheres From Very Low-Mass Stars to Extrasolar Planets"
- Alvarez R., Plez B., 1998 A&A 330, 1109
 "Near-infrared narrow-band photometry of M-giant and Mira stars: models meet observations"
- 3. Asplund, M., Grevesse, N., Sauval, A. J., Scott, P. 2009, ARA&A 47, 481–522 "The Chemical Composition of the Sun"
- Baraffe, I., Homeier, D., Allard, F., Chabrier, G., 2015, A&A 577, A42 "New evolutionary models for pre-main sequence and main sequence low-mass stars down to the hydrogen-burning limit"
- Branco, V., Coelho, P. R. T., Lançon, A., Martins, L. P., Prugniel, P. 2024.
 "Synthetic stellar spectra to study multiple populations in globular clusters: an extended grid and the effects on the integrated light." arXiv e-prints. doi:10.48550/arXiv.2404.15468
- 6. Caffau, E., Ludwig, H.-G., Steffen, M., Freytag, B., Bonifacio, P., 2011, Solar Physics 268, 255–269
 "Solar Chemical Abundances Determined with a CO5BOLD 3D Model Atmosphere"
- Chiavassa, A., Casagrande, L.; Collet, R.; Magic, Z.; Bigot, L.; Thevenin, F.; Asplund, M., 2018, A&A 611, A11
 "The Stagger-grid: A grid of 3D stellar atmosphere models V. Synthetic stellar spectra and broad-band photometry"
- De Laverny P., Recio-Blanco A., Worley C.C. and Plez B., 2012 A&A ,544, A126 "The AMBRE project: A new synthetic grid of high-resolution FGKM stellar spectra"
- 9. Grevesse N. and Sauval J., 1998, SSRv 85, 161 "Standard Solar Composition"
- 10. Gustafsson B., Bell R.A., Eriksson K., Nordlund Å., 1975 A&A 42, 407 "A grid of model atmospheres for metal-deficient giant stars. I"
- Gustafsson B., Edvarsson B., Eriksson K., et al., 2003 in "Stellar Atmosphere Modeling", ASP Conf. Ser. Vol. 288, p331 "A Grid of Model Atmospheres for Cool Stars"
- Hillier D.J., Miller D.L., 1998 ApJ 496, 407
 "The Treatment of Non-LTE Line Blanketing in Spherically Expanding Outflows"
- 13. Hubeny, I. & Lanz, T., 2000, http://nova.astro.umd.edu/Tlusty2002/pdf/syn43guide.pdf
- 14. Kupka F.G., Ryabchikova T.A., Piskunov N.E. et al., 2000 Baltic Astronomy 9, 590 "VALD-2 – The New Vienna Atomic Line Database"

- Kurucz R.L., 1993, IAU Coll. 138, in "Peculiar versus normal phenomena in A-type and related stars" ASP Conf. Ser., Vol. 44, p.87 "A New Opacity-Sampling Model Atmosphere Program for Arbitrary Abundances"
- 16. Kurucz R.L., 2005, Memorie della Societa Astronomica Italiana Supp. v8., 14, "ATLAS12, SYNTHE, ATLAS9, WIDTH9, et cetera"
- Magic, Z. and 7 colleagues, 2013, A& A 557, A26
 " The Stagger-grid: A grid of 3D stellar atmosphere models. I. Methods and general properties"
- 18. Marcolino, W., Bouret, J.-C., Martins, F., Hillier, J., 2024, submitted to A& A
- Meynet G. and Maeder A., 2003, A&A 404, 975
 "Stellar evolution with rotation. X. Wolf-Rayet star populations at solar metallicity"
- 20. Palacios A., Gebran M., Josselin E., Martins F., Plez B., Belmas M., Lèbre A., 2010, A&A 516, A13
 "POLLUX : a database of synthetic stellar spectra"
- Plez B., Brett J., Nordlund Å., 1992, A&A 256, 551
 "Spherical opacity sampling model atmospheres for M-giants. I Techniques, data and discussion"
- 22. Schaller G., Schaerer D., Meynet G., & Maeder A., 1992, A&AS 96, 269 "New grids of stellar models from 0.8 to 120 solar masses at Z = 0.020 and Z = 0.001"
- 23. Seaton M. J., 2005, MNRAS, 362, L1 "Opacity Project data on CD for mean opacities and radiative accelerations"
- Badnell N. R., Bautista M. A., Butler K., Delahaye F., Mendoza C., Palmeri P., Zeippen C. J., Seaton M. J., 2005, MNRAS, 360, 458 "Up-dated opacities from the Opacity Project"
- 25. Gray D. F., " The observation and analysis of stellar photospheres", Cambridge University Press, Third Edition